



Towards Design Guidelines for Effective Health-Related Data Videos: An Empirical Investigation of Affect, Personality, and Video Content

Samar Sallam
University of Manitoba
Winnipeg, Canada
sallams@myumanitoba.ca

Yumiko Sakamoto
University of Manitoba
Winnipeg, Canada
umsakamo@umanitoba.ca

Jason Leboe-McGowan
University of Manitoba
Winnipeg, Canada
jason.Leboe-McGowan@umanitoba.ca

Celine Latulipe
University of Manitoba
Winnipeg, Canada
celine.latulipe@umanitoba.ca

Pourang Irani
University of British Columbia
Kelowna, Canada
pourang.irani@ubc.ca




Figure 1: Data Videos, or animated infographics, are short videos that present large amounts of data in an easy and engaging narrative format. For Study 1, we selected nine Data Videos focusing on three health-related topics: physical activity, sleep, and diet. This figure shows an example of the videos used in the study. The full list of videos and more detailed descriptions are listed in Appendices A & B.

ABSTRACT

Data Videos (DVs), or animated infographics that tell stories with data, are becoming increasingly popular. Despite their potential to induce attitude change, little is explored about how to produce effective DVs. This paper describes two studies that explored factors linked to the potential of health DVs to improve viewers' behavioural change intentions. We investigated: 1) how viewers' affect is linked to their behavioural change intentions; 2) how these affect are linked to the viewers' personality traits; 3) which attributes of DVs are linked to their persuasive potential. Results from both studies indicated that viewers' negative affect lowered their behavioural change intentions. Individuals with higher neuroticism exhibited higher negative affect and were harder to convince. Finally, Study 2 proved that providing any solutions to the health problem, presented in the DV, made the viewers perceive the videos as more actionable while lowering their negative affect, and importantly, induced higher behavioural change intentions.

CCS CONCEPTS

• **Human-centered computing** → **Human computer interaction (HCI); Empirical studies in HCI.**

KEYWORDS

Data Video, narrative visualization, data storytelling, persuasive technology, attitude change, personality, affect, actionable, solution, physical activity

ACM Reference Format:

Samar Sallam, Yumiko Sakamoto, Jason Leboe-McGowan, Celine Latulipe, and Pourang Irani. 2022. Towards Design Guidelines for Effective Health-Related Data Videos: An Empirical Investigation of Affect, Personality, and Video Content. In *CHI Conference on Human Factors in Computing Systems (CHI '22)*, April 29-May 5, 2022, New Orleans, LA, USA. ACM, New York, NY, USA, 22 pages. <https://doi.org/10.1145/3491102.3517727>

1 INTRODUCTION

Videos consisting of motion graphics with visual and auditory stimuli to tell a *story centered around empirical data*, called Data Videos (DVs) [5], are becoming an increasingly popular communication medium, often aimed at promoting behavioural change. For instance, a 2020 study examining YouTube videos covering COVID-19 vaccine development asserts that such videos are considered one of the major sources of information for many people [12]. While this popularity could be attributed to technology advancement (e.g., DVs can be viewed with smartphones and tablets anytime and anywhere), another reason could be attributed to DVs' ability to attract broad audiences on any given topic due to their narratives [37]. DVs can raise viewers' awareness about social issues, be

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.
CHI '22, April 29-May 5, 2022, New Orleans, LA, USA

used for advertisement, and more recently for promoting healthy behaviours [20] for a wide range of viewers regardless of their data literacy background. Furthermore, DVs are short [5, 41], and obtaining information quickly is critical in today's fast-paced lifestyle. Additionally, DVs can present *actionable intelligence* components. A majority of COVID-19-related DVs, for example, focus on social distancing, mask-wearing, and vaccination, which are all actions that can be taken to reduce the spread of the disease. These videos usually show statistics and facts with a story that plays on viewers' affect, while drawing data-driven conclusions to encourage smart decisions. In short, DVs provide data-driven *actionable insights*.

Data visualization, or finding ways to best present the data at hand, is a major interest for HCI researchers. This interest is growing in parallel with the recent era of big data as enormous amounts of data are becoming available due to the technological advancements in sensors and embedded systems (e.g., smartphones). This naturally leads us to focus more on the visualization aspect in order to present the data in the most insightful and understandable way. Storytelling and narratives, as a natural means for human communication, have attracted the attention of visualization researchers and have proven to be very effective [37]. While visual data storytelling that depends on pictures and graphs (i.e., infographics), has proven efficacy in the field of data visualization, Gershon and Page argue that putting such visuals in a video format can take the story to a whole new level [37]. While "a picture is worth a thousand words", a video that elaborates and supports that picture with animation, audio narration, music, and sound effects is considered to be "worth a million pictures" according to Gershon and Page [37]. The potential of videos for data storytelling has been further amplified with the emerging popularity of data-based videos, or DVs, on social media platforms such as Youtube, Facebook, Instagram, and TikTok, which are easily accessible to a wide audience using various smart devices.

While DVs are beginning to attract the attention of HCI and visualization researchers to understand their building blocks [5] and provide tools to create such videos [7, 60], they are still a novel form of data visualization [5, 6, 20], and human perception in the context of DVs has not been explored fully yet despite its relevance. In this research, we explore DVs targeting common health issues (physical activity, sleep, and diet) to understand the factors affecting their ability to influence their viewers' health-related behavioural intentions. Given the brevity of DVs [5, 41], we focus on DVs' ability to influence their viewers' perception of the topic (i.e., targeting viewers' intentions/willingness to improve their behaviour, but not actual behavioural change). We chose to focus on generic behavioural health issues as opposed to specific health issues (e.g., diabetes or insomnia) for two reasons. First, due to the higher relevance of those general topics to the majority of the population, and hence relevance to our randomly recruited participants. Further, we believe that people with serious health issues are probably dealing with their issues with their own doctors (unique advice might be needed). While simple and well known to everyone, common health-related behavioural issues, such as poor eating habits and physical inactivity, could be linked to serious chronic consequences (e.g., cardiovascular diseases, obesity, and even some types of cancer) [1, 64]. People suffering from behaviour-induced health issues could benefit if they understood how their

seemingly minor behaviours underlie their serious medical issues, or alternatively, the potentially beneficial outcomes if they changed those behaviours. We believe that the videos could serve as good reminders in the context of our study. DVs have a great potential to help in this area because of their compelling narrative nature. The storytelling aspect of DVs often arouses emotions in the viewers' minds. Research in advertisement and marketing demonstrates that the extent of such emotional experiences, or *affect*, plays a significant role in motivating viewers [45, 51, 56]. For example, one starts exercising after *feeling bad or at risk*. It is important to note that personality differences could underlie individuals' unique affective responses to a persuasive message. Indeed, individuals' responses to an affect-provoking persuasive message are unique [42, 46, 88]. Thus, investigation of viewers' personality traits could lead us to fruitful DV personalization strategies. Our exploration is crucial in identifying basic approaches to design DVs, as a data representation interface, from an HCI perspective to efficiently influence their viewers in a short amount of time. Understanding such personalization strategies has the potential to lead to new DV design tools that assist designers in making more effective and impactful DVs.

To explore the effect of DVs on viewers' perception, Study 1 focused on three questions: **RQ1**: how is viewers' affect, induced by DV viewing, linked to their willingness to alter their health-related behaviours?; **RQ2**: how are these affective responses linked to the viewers' personality traits?; and **RQ3**: what are the underlying attributes in DVs that contribute to their persuasive potential? Subsequently, to provide guidelines for video designers, Study 2 explored how we can tune participants' affect along with the narrative structure, with *useful information*, to improve the overall impact of DVs by altering the types of information.

2 LITERATURE REVIEW

First, we describe research investigating the effectiveness of Data Videos (DVs) as a narrative to communicate data, as this is the foundational media our work explores. We highlight some behavioural theories and studies in HCI and other related fields because we are interested in how DVs can promote positive behavioural changes. We then discuss work related to how affect is impacted by narrative structures, and how it could contribute to behavioural change intentions. As a factor influencing affect, we review studies about how personality traits impact responses to persuasive messages, by describing studies and strategies used in the literature of persuasive technologies. Finally, we briefly describe studies exploring measurements to assess the effectiveness and quality of health-related videos.

2.1 Data Videos as a Narrative

Data Videos (DVs) are motion graphics that incorporate factual, data-driven information to briefly tell informative and engaging stories with data [5-7], typically shorter than six minutes [41]. DVs are gaining popularity in various fields such as journalism [89, 90], education [41, 85], advertising and mass communication [78], as well as in political messages and campaigns [32, 79]. Due to their narrative nature, DVs are recognized as one of the seven forms of narrative visualizations [18, 77]. The power of DVs comes mainly from this narrative format; stories can convey information in an

engaging way that is more natural, seamless, and effective than text or even pictures [37, 43]. A well-told story can convey a large amount of information in a short period of time, in a way that viewers find interesting and easy to understand [37]. Moreover, they tend to be more believable and memorable [37, 48].

Narrative Transportation Theory states videos can transform and immerse the viewer in a totally different world [87]. The locations, characters, and emotion-laden scenarios then generate the opportunity to impact the viewers' own beliefs, emotions, and intentions [40, 60, 61]. Furthermore, a solid wealth of psychological theories support the persuasive power of narrative. The Extended Elaboration Likelihood Model argues that as people indulge in the narrative, with all its cues and stimuli, their cognitive processing of the narrative obstructs making counterarguments in response to the presented message [17, 81], making the message more persuasive even for those who are difficult to persuade otherwise [80]. Additionally, as per the Entertainment Overcoming Resistance Model, the entertaining aspect of a narrative also plays a role in reducing the cognitive resistance to the message presented, and hence facilitates persuasion [22, 59, 60].

Although DVs have great potential for motivating changes in behaviour, and have become an increasingly relied upon source for accessing information, it was not until recently that empirical investigations have focused on the component features and narrative characteristics that help to determine their capacity to persuade [5, 20]. Baber et al. [9] define narrative as a formal structure that constitutes a "shareable" story or report as opposed to the informal stories which could be "unstructured" and "ambiguous". The narrative is defined as a series of connected events that constitute a story [77]. The order in which these events are presented in a DV constitutes its narrative structure, and influences the understanding and the viewing experience of the video [23]. Amini et al. [5] examined 50 professionally created DVs to learn about their narrative structure. In their study, they divided the videos into temporal sections and coded them based on Cohn's [23] theory of visual narrative structure that broke the narrative into four categories: Establisher, Initial, Peak and Release. The Establisher is the entry point to the video which introduces its context, the Initial initiates the action, the Peak is where the most important events and the tension occurs, and the Release is where the conclusions are drawn [5, 23]. After analyzing and coding the videos based on these categories, the authors found that these categories are ordered differently based on video types: factual videos with multiple facts and information; comic and open-ended videos that do not have a conclusion or a takeaway message; and videos that exhibit multiple problems or those that provide multiple solutions to a single problem. The authors pinpointed some narrative structure patterns that are commonly used in DVs. By far, the most commonly used structure was the "well-balanced" pattern that strongly introduces and establishes the stories with facts till the peak is reached, then takes a descending direction towards the solutions or takeaway messages in the Release. The second most common pattern is one that ends at the peak leaving the viewers with an open question or unresolved problem.

Choe et al. [20] introduced a new class of Data Videos called Persuasive Data Videos or PDVs. This genre of DVs incorporates

some persuasive elements inspired by and drawn from Oinas-Kukkonen and Harjumaa's [65] Persuasive System Design Model. Choe et al. [20] studied how incorporating some persuasive elements in a DV could improve persuasive potential of the video. Their PDVs had higher persuasive potential than regular DVs. Amini et al. [6] examined the effect of using pictographs and animation, two commonly used techniques in DVs. They found that the use of such techniques enhanced the viewers' understanding of data insights while boosting their engagement.

2.2 Affect and Data Video

In their investigation of pictographs and animation in DVs, Amini et al. [6] concluded that pictographs are effective because they trigger more emotions in their viewers, while the animation intensifies such emotions. This leads us to an important aspect of DVs: the viewers' emotional experience, or *affect*. Viewers' preference for multimedia, be it a performing art, internet video, or even music videos, is highly dependent on their arousal level and the intensity of their affect towards the media [10, 83]. Early studies assumed that TV viewers liked to watch shows that elicit positive emotions as opposed to negative emotions [38]. However, later research showed that while this might be true for real life events, it was not true for TV shows [62]. In a more recent study, Bardzell et al. [10] examined the intensity and valence of viewers' affect, as well as their ratings of internet videos. The results showed a correlation between the affect intensity and the liking of the video. As for the valence (i.e., positive or negative), the study showed that it is not the presence or absence of negative or positive affect that influenced the rating of the video. Rather it is the *emotional arc* that leaves the viewer emotionally resolved and hence liking the video, even if it started with negative emotions. This shows that negative emotions should be resolved in order to generate a satisfactory user experience with a narrative.

2.2.1 The Role of Affect in Attitude and Behaviour Change.

Affect plays an important role in the appeal as well as the persuasive power of media content [2, 15, 51]. According to theories in psychology, some of our attitudes have a cognitive base while others have an affective base [47, 69]. *Affective attitudes* emerge from our feelings towards certain topics or ideas. Some attitudes are influenced relatively easily through affect while others through logic and thoughts (i.e., cognition) [47, 69]. The Dual Process Model suggests two routes to persuasion: central and peripheral. The central route is the cognitive route in which the receiver of a message is willing and able to cognitively process the ideas [17, 69]. In contrast, the peripheral route processing is taken when the receiver lacks the motivation or ability to logically process cues in the message, and decides to agree with the message based on its emotional appeal (e.g., emotions triggered by the look or smell but not by the logic) [17, 51]. For instance, one might purchase a car based on its gas emission, cost, or functions (central route) or because of the way it looks (peripheral route). Thus, both cognition and affect are heavily involved in persuasion. Data Videos constitute a perfect persuasive medium that combines both the cognitive aspects (fulfilled by the data), and the emotional aspects (fulfilled via the narrative and the supporting audio and visual elements) needed for the persuasion process. While there has been some empirical

investigation of what types of data visualizations should be used in DVs [5], no clear guidelines have been provided regarding the effective emotional messaging strategies and techniques that ought to be used when developing DVs that aim at influencing the viewers' perception about health-related behaviours. In the field of marketing, an area focusing primarily on persuading and guiding the viewers to adopt a certain service or commodity, a wide array of studies have examined the kind of affect evoked by advertisements [45] and how they influence the viewers' attitudes [16]. However, no research has explored the function of affective changes induced by Data Videos as a potential persuasive medium.

2.2.2 *Affect Valence and Persuasion.*

When it comes to health messaging aiming at attitude or behaviour change, the framing of the message often takes one of two strategies: a gain-framed strategy that focuses on the gains or positive consequences of following the healthy behaviour (i.e., positive emotions such as motivation or happiness); or a loss-framed strategy that focuses on the losses or the negative outcomes of failing to follow the healthy behaviour (i.e., negative emotions such as fear or worry) [34]. The Prospect Theory suggests that the same information can have different effects on their receivers based on how they are framed [34, 86]. The theory suggests that people would be more eager to avoid potential losses than they would be to attain potential gains. Rothman and Salovey [74] suggest the use of gain-framed strategy for persuasive messages targeting illness prevention behaviours such as physical activity, while a loss-framed strategy would be more effective for messages targeting illness detection behaviours like mammography [34]. The model of behaviour change, on the other hand, suggests that negative affect such as feeling worried or at risk is the first step towards attitude change [40]. In short, some research found significant effects of positive affect on behaviour change [68], some found undesirable effects of using negative affect [92], while others recommend using a threatening or fear-inducing approach for health issues such as smoke cessation [67] and physical activity [14]. While both approaches are worth exploring, in our studies we decided to focus on and explore the effects of using negative affect due to the nature of the existing high-quality videos chosen for the study based on the researchers' consensus.

2.2.3 *Measuring Affect.*

There are two approaches to measuring affect: 1) the implicit approach, which relies on physiological recordings of individuals' biometric responses; and 2) the explicit approach, which relies on self-reporting of the users' affect during their exposure to the stimuli. While sensors and devices that can explicitly log biometric changes related to affect are very promising (e.g., heart rate, breath rate, respiration patterns, and galvanic skin responses (GSR)) [50, 83], they are often invasive, expensive, and the meaning of the data recorded can be ambiguous [50]. As for the explicit measurement methods, indicators such as final applause to a show, post-show surveys, interviews, as well as different scales are most commonly used. While less costly, such explicit approaches can capture somewhat skewed responses as the viewers' responses are normally affected by their peak emotion and the emotions experienced by the end of the show (i.e., the 'peak-end' effect) [26, 49, 50, 72]. Latulipe et al. [50] measured participants' GSR to a video in relation to their

responses to two self-report scales. They found correlations between the explicit and implicit measures of emotions, showing that self-reporting of emotions can be reliable. Due to the online nature of our studies that was enforced by the pandemic, we adopted the *explicit* approach using scale items based on the Spielberger State Anxiety Scale [55].

2.3 Personalization in persuasive technology

Recent research indicates that the one size fits all model of persuasive technology is not so effective. Instead, researchers are showing the power of *personalized* persuasive systems which often explore the effect of personalities on persuasion [8, 19, 27, 44, 46, 47, 88, 91], making it essential for us to consider the personality aspect as a factor in our studies. The five-factor model (FFM) of personality offers five broad traits, referred to as the "Big Five", that capture differences in typical patterns of thought and behaviour that vary across individuals: Extraversion, Agreeableness, Conscientiousness, Neuroticism, and Openness to Experience. The FFM is the most widely accepted model for personality assessment across diverse disciplines [29, 36, 47, 57, 76, 88] hence, we adopted it for our personality assessment. Halko and Kientz [42] explored the link between personality traits and people's perception regarding different persuasive strategies. They found correlations between participants' personality traits and persuasive strategies. For example, people who score high in Neuroticism tend to prefer negative reinforcement (i.e., removal of aversive stimuli), as opposed to positive reinforcement (i.e., rewards) to improve their behavioural patterns. They also found that neurotic people do not like cooperating with others to achieve their goals. Loss-framed strategies are commonly used in health DVs, so in our studies, we investigate the corresponding effects of this strategy on the viewers, given their personality differences.

2.4 Measures for assessing health-related videos

The content of a video, with its particular visual/auditory elements, animations, graphics, narration style, etc., contributes to the video's potential to affect attitude/behavioural change intentions. Various studies have examined ways to assess the quality of YouTube videos tackling health-related issues like exercise and low-fat diets [13, 31], bruxism [35], or general health videos [28, 33, 75]. For assessing the quality of YouTube Videos addressing health issues, Gabarron et al. [33] examined: 1) the usefulness and quality of the content; 2) Popularity-Driven Measures (i.e., number of views, shares, and video rating); 3) Heuristic-Driven Measures (e.g., length, narrative, visuals). However, quality assessment of such videos has been mainly done by experts (e.g., health professionals, major health organizations, IT researchers) [33] and very few studies have considered the feedback or judgements of the patients or viewers of such videos [24, 33].

In sum, research supports the potential of DVs to effectively convey health-related data through their affective influences. Studying affective responses in conjunction with personality traits is appealing because, understandably, affective responses are guided by individuals' traits. Exploration of personality traits and affective responses will show us ways to effectively personalize DVs for

better communication. Additionally, to investigate common video attributes from the viewers' perspective, in relation to potential attitude change, will be beneficial. While previous studies in other fields (e.g., psychology) explored related constructs [2, 15, 51]), our study focused on the users' perception in the frame of Data Videos in an applied manner so the designers of DVs can benefit from our research: Our explorations would contribute in providing general guidelines for DV designers who are interested in presenting health information.

3 STUDY 1: EXPLORATION OF AFFECT, PERSONALITY, AND POTENTIAL FOR ATTITUDE CHANGE

To understand the underlying mechanisms of health-related Data Videos (DVs) in relation to viewers' experienced affect, their personality traits, as well as the persuasive potential of DVs, Study 1 aimed to answer three main questions: 1) Does increasing viewers' negative affect levels lead to higher potential persuasion?; 2) Is there a relationship between participants' personality traits and their affective state after watching DVs?; and 3) What predicts higher potential attitude change in terms of personality traits as well as video content?

3.1 Study Administration

An online study was created using Qualtrics, and administered through Amazon Mechanical Turk (MTurk). To ensure higher data quality, we recruited MTurk workers who had completed a minimum of 1000 tasks prior to our study, with approval ratings higher than 95%. All participants received monetary compensation (\$2.24 US) in compliance with the study ethics approval and MTurk payment terms. US and Canadian participants who were older than 18 were recruited.

3.2 Data Video Selection

We collected DVs focusing on three general health-related topics: physical activity, healthy sleep, and healthy diet. We systematically explored existing DVs with guidance taken from Amini et al.'s [5] study.

- First, two researchers collected over 100 videos using keywords such as 'healthy diet', 'dangers of not having enough sleep', 'importance of exercise', etc.
- We then removed videos containing erroneous information and/or did not follow Amini et al.'s [5] DV definition.
- Two researchers coded the remaining videos for *length*, *view count*, *source credibility*, and *information accuracy*.

The final list consisted of nine videos; three on each topic (see Appendix A).

3.3 Data Collection Instruments

Personality Traits We utilized a shortened, 10-item, version of the Big-Five Inventory [71] (e.g., "I see myself as someone who gets nervous easily", using a 5-point Likert scale) to assess participants' personality traits. This scale is widely accepted in personalized technology research [8, 66] in which personality assessment needs to be quick, and has been repeatedly validated. For instance, Gosling

Table 1: General Persuasive Potential (GPP) items of the Persuasive Potential Questionnaire (PPQ)

I feel that...

- the video would make its viewer change their behaviours.
 - the video has the potential to influence its viewer.
 - the video gives the viewer a new behavioural guideline.
-

A Likert scale (1: Strongly Disagree to 7: Strongly Agree) is used.

et al. [39] noted it has "reached an adequate level" in terms of predictive power and convergence with full scales in self, observer, and peer responses.¹

Affective State Self-Reports Participants' negative affect were assessed using 3-items: "I feel anxious"; "I am relaxed (R)"²; and "I am worried", using an 8-point Likert scale based on the Spielberger State Anxiety Scale [55] where they eliminated a neutral choice. Out of the final nine videos we found suitable for the study due to their credibility, quality, and information accuracy, seven videos followed a loss-framed messaging strategy, and hence we focused on negative affect.

Persuasive Potential Questionnaire (PPQ) While DVs might have a great potential to influence their viewers, they are brief video clips. Hence, we suspect only one exposure to a video (as in our study) might not induce any effect at the behavioural level. Therefore, we focused on DVs' *potential* for persuasion at the perceptual level. To measure the potential of DVs, the Persuasive Potential Questionnaire (PPQ) [58] was adopted and adjusted to fit our context. We focused on the General Persuasive Potential (GPP) dimension of PPQ, see Table 1.

Video Content Appraisal Three questions examined participants' *appraisal of the video content* focusing on information characteristics: Novelty, Clarity, and Usefulness of the information in each DV (e.g., "The information provided by the video was useful to me", using a 7-point Likert scale, 1: Strongly Disagree to 7: Strongly Agree).

3.4 Study Progression

First, participants answered demographic questions, followed by the personality measure and the affect questions (i.e., baseline affective state). Participants then watched three DVs; one video from each topic, and answered questions after each. The videos were randomly selected from the sets of three videos per topic. The topic order was also randomized to avoid any priming and/or order effects. After watching each video (three videos in total), participants responded to:

- (1) The three Affect-related items, for a post video measure;
- (2) Video content appraisal questions;
- (3) Questions for the General Persuasive Potential (GPP) of the video (see Table 1);
- (4) Finally, participants reported any health issues that would prevent them from following the video's advice.

¹According to Google Scholar search, this scale has been cited in 3468 articles at the moment of writing; August, 2021

²(R): item is reverse scored

After completing these four steps for each video, participants engaged in a 12-piece jigsaw puzzle game, which was designed to take at least one minute. This step was added based on various psychology studies in which a filler task (e.g., a visuospatial task like a word puzzle, counting numbers from 7 to 0) is used between tasks [11, 54, 82] to help participants neutralize their affective state by focusing on an unrelated task [70, 82]. After the filler task, participants repeated the four steps for the next video.

3.5 Results

On average, participants took 26 minutes to complete the study. Data-fitting assumptions for each analysis were checked and non-parametric options were used with report of medians instead of means, whenever appropriate. Due to our smaller sample size, more detailed results including null effects along with effect sizes (which is independent from sample size issues [84]) are reported in Appendix F so the readers can examine the potential Type II error.

3.5.1 Participants.

One hundred and two people with ages ranging between 21 and 70 participated in our study ($M = 37.29$, $SD = 12.01$): 68 identified as male, 33 as female, and one preferred not to report; 61% of them self-identified as white, 21% did not report their ethnicity, and the rest self-reported as Hispanic, Black, or Asian. Almost all of the participants (98%) reported their first language was English, and 83.3% of them reported they had at least a Bachelor's Degree.

3.5.2 Data Quality Control.

To filter out potentially unreliable responses, a verifiable (i.e., Gotcha) question was designed to be readily solvable as long as the participants read the question ("How many words do you see in this sentence?"): 78 cases remained for the analyses³. When appropriate, we further filtered out responses based on their answer to a question ("I have health issues that prevent me from following the advice provided in the video") in each of the three topics. This was designed to ease potential confounds as much as possible (i.e., participants' lack of willingness to change their behaviour might not be reflecting their reactions to the DVs. Instead, this could be due to their health issues)⁴.

3.5.3 Data Preparation.

First, we explored inter-item reliability for each scale (e.g., GPP, affective response, etc.), and when Cronbach's Alphas were higher than our standard ($\alpha \geq .70$) based on Nunnally and Bernstein [63], we computed the mean across all items to be used as an index:

- **Negative Affect Index:** Cronbach's Alphas were higher than our standard ($.73 \leq \alpha \leq .82$) for participants' negative affective responses per topic (Physical Activity, Sleep, and Diet), indicating that items for each scale had sufficient internal consistency.⁵ Thus, the mean of three items was computed for each topic. Then the correlations between these means for each topic were also investigated. They were all significantly

and highly correlated ($.810 < r_{hos} < .830$, $ps < .001$) [4]), implying that if a participant's affective response was negative from viewing one video, it was likely that they experienced negative affects from viewing other videos as well (i.e., implied underlying *personal tendency*). Thus, the mean across all the topics was used as an index for Negative Affect.⁶

- **GPP Index:** The index for GPP was also created in the same manner. Cronbach's alphas ranged between .81 and .91 per topic. Further, GPP for one topic was correlated with the GPP for other topics ($.555 < r_{hos} < .719$, $ps < .001$)⁷, and the mean of scores across all the topics was generated, and used as a GPP index.⁸

3.5.4 Data Analysis and Findings.

Negative Affect and GPP: We explored whether overall GPP could be predicted by negative affect. Negative affect predicted GPP, $F(1, 76) = 4.056$, $p = .048$, $R^2_{adj} = .038$, $\beta = -.225$. Interestingly, participants' increased levels of negative affect predicted lower GPP, (i.e., it was harder to motivate the viewers to alter their health-related behaviour when they had high levels of negative affect after viewing the video).

Predictors of GPP: We conducted a multiple regression analysis, using a stepwise method, to explore how content appraisal variables (i.e., Information Novelty, Clarity, and Usefulness), and personality traits could predict GPP altogether. Content appraisal and personality traits are unique as content belongs to the DV design while personality traits belong to individual viewer. Thus, they are conceptually independent routes to influence the viewers' perception (design itself & target selection). Only two predictors remained in the model, which explained approximately 66% of the variability in GPP (1; Information Usefulness, $\beta = .718$, $p < .001$, 2; Neuroticism, $\beta = -.143$, $p = .036$), $F(2, 75) = 74.87$, $p < .001$, $R^2_{adj} = .657$ (see Fig 11 in Appendix F for details on the excluded variables). When viewers perceive the information in DVs as useful, it was particularly easier to influence the viewers' attitude. However, DVs will be less effective at convincing individuals who have highly neurotic tendency to change health-related behaviours. For those who score high on neuroticism, a differently framed narrative may be more effective. Our model confirms that consideration of both routes (i.e., video design & target selection) are useful in achieving higher persuasive potential.

Personality Traits and Negative Affect: To examine the relationship between personality traits and the levels of negative affect, correlations between each personality trait and negative affect were explored (see Fig 12 in Appendix F). Negative affect was positively correlated with neuroticism, $\rho(78) = .594$, $p < .001$, but negatively correlated with conscientiousness, $\rho(78) = -.363$, $p = .001$; no other traits were correlated with negative affect (b in Figure 2). Highly neurotic people experience higher levels of negative affect by DV viewing, unlike conscientious people who are less sensitive. **Data Videos and Negative Affect:** Next, we turned to examine the characteristics of DVs. As seen in Figure 3, each DV induced

³23.5% of our participants failed to pass our Gotcha question.

⁴Four participants responded "Yes" after viewing a physical activity themed video, four others responded "Yes" after watching a sleep themed video, and finally, three others responded "Yes" after watching a diet themed video. A pairwise deletion method was applied to this selection throughout the analyses.

⁵One item ("I am relaxed") was reverse coded for the index computation.

⁶Negative Affect refers to the Negative Affect Index wherever mentioned.

⁷This implied another underlying *personal tendency*; if a participant perceived high levels of general persuasiveness in one video, they perceived higher levels of general persuasiveness in other videos as well

⁸Mentions of GPP in the analysis refer to the overall GPP Index.




Figure 2: Negative Affect was correlated with the Video Content (a) and Personality Traits of Neuroticism and Conscientiousness (b). Finally, Negative Affect predicted viewers’ unwillingness to alter their behaviours (c).

*: negative correlation

different levels of negative affect. DVs that induced the highest (Diet 1) and the lowest (PA 1) negative affect were examined closely. Wilcoxon Signed Ranks Tests revealed that viewers’ negative affective responses were increased after watching Diet 1 Video (Pre-video $Mdn = 3.00$; Post-video $Mdn = 5.00$, $p = .001$), but not after the PA1 Video (Pre-video $Mdn = 2.00$; Post-video $Mdn = 1.67$, $p = .48$). To understand this differential effect, two researchers coded the videos’ contents qualitatively and found that Diet 1 Video included a fear inducing message without any clear solution. Consequently, participants’ negative affect remained high even after the video. We further observed that none of the three DVs that induced the highest levels of negative affect (see Fig 3) contained concrete solutions (see Appendix B). Thus, it appeared that not providing solutions was linked to heightened negative affect.




Figure 3: Negative Affect by video. (Note PA = Physical Activity). The horizontal dotted line indicates the mean across all 9 videos ($M = 2.96$). Videos that induced the highest negative affect (Diet 1, PA 3 & Sleep 1) did not have solutions.

3.6 Discussion

Due to the nature of common health Data Videos which regularly contain fear-inducing messages, this study explored negative affect (i.e., anxiety, worries, and not being relaxed) in response to those messages specifically about physical activity, sleep, and diet. We observed that DVs without clear solutions to their fear-inducing messages kept viewers’ negative affect higher. We also found that heightened levels of negative affect predicted a lower viewer willingness to change their behaviour. Moreover, the levels of negative affect viewers experienced were linked to their own personality traits: individuals who scored high on neuroticism experienced higher levels of negative affect while those who scored high on conscientiousness were less likely to experience highly negative affect. These findings flag the potential risk of including heavily loss-framed messages (i.e., messages that focus on negative outcomes of not following a certain behaviour) in DVs especially for highly neurotic individuals who have higher tendency to experience negative affect while being harder to persuade. While loss-framed messages are commonly included in DVs, and their effect is validated in psychological research, such messages could potentially backfire in DVs: the messages could be perceived as less convincing when a suitable or useful solution is not presented. Finally, perceived information usefulness played an important role in predicting GPP. However, information usefulness is a broad construct: understanding what constitutes information usefulness might be essential in producing powerful DVs. We acknowledge that our sample size became smaller than we originally planned, after removing invalid data points, and another study with a larger sample size will be fruitful especially in examining our null effects (see Appendix F). We made several important discoveries (e.g., how information usefulness is related to GPP, how negative affect influences GPP within DVs) even with the smaller sample size, however. Thus, Study 2 further explores how we can structurally resolve negative affect with useful information.

4 STUDY 2: EXPLORATION OF THE EFFECT OF PROVIDING SOLUTIONS TO HEALTH-RELATED DVs ON THE LEVELS OF NEGATIVE AFFECT AND PERCEIVED USEFULNESS.

To follow up on our findings in Study 1 regarding perceived information usefulness, we developed a nomological network to break down the construct (see Appendix C). The overall goal of Study 2 was to explore the means to effectively resolve the negative affect along the narrative arc with *useful* information. Our literature review revealed that an actionable solution is a part of perceived information usefulness (Appendix C). Interestingly, since a lack of solution was linked to viewers’ negative affect in Study 1, we decided to explore the role of solutions in DVs. Specifically, we investigated the effect of providing two types of solutions: 1) a broad/generalizable solution, and 2) a concrete/actionable solution, compared to having no solution at all (baseline). We hypothesized that, by providing concrete solutions, the viewers’ anxiety levels could be assuaged as they feel the issues are resolvable. This hypothesis is based on results from Study 1 and Cohn’s [23] narrative

structure that starts with introducing the problem (*Establisher*), followed by a rise in action (*Initial*) where the events intensify, or the negative consequences of not following a healthy behaviour (in the case of DVs targeting health issues) are listed, till the *Peak* is reached. Finally, the narrative takes a descending direction towards a resolution to the problem (*Release*). This ‘well-balanced’ narrative structure is very common in DVs [5] and is consistent with Bardzell et al.’s [10] findings regarding viewers’ liking of general internet videos. They found that viewers liked videos that succeeded to utilize an *emotional arc* that managed to resolve the issues discussed by the end of the video, even if the video started with negative emotions [10]. Hence, we anticipated that presenting the issue will increase the level of anxiety, but this will be soon alleviated if solutions are presented. This reduction in anxiety level might be a key to improving the persuasive potential of the videos. We also believe that the effect of providing a solution is twofold: it should reduce viewers’ negative affect, and should improve the perceived usefulness of the video which then aids in the persuasive potential.

4.1 Study Administration

We chose physical exercise as a topic because since the COVID-19 pandemic started, many people lost their access to the gym or were prevented from participating in any training or group physical activities (i.e., reduced physical activity level). Parallel to Study 1, Qualtrics survey was administered through MTurk with the same workers’ requirement criteria with monetary compensation (\$2.47 US). Again, participants were from the US and Canada. The study was a between-subjects design with three conditions: Control/No Solution, Generalizable Solution, and Concrete Solution. Participants completed survey questions before, after, and between segments of a DV on physical activity.

4.2 Data Video Development

We used a DV developed by the World Health Organization, titled ‘Move for Health’, as the basis for our stimuli [30]. The original video starts by stating that modern life has made us less physically active (*Establisher* in the narrative arc (see the yellow arc in Fig 4). The video lists the dangers of physical inactivity in the *Initial* phase (e.g., fourth leading risk factor for mortality worldwide, a major cause of severe diseases such as diabetes, etc.), the video then reaches its *Peak* by referring to office workers who remain seated for eight hours a day and how that could lead to severe health issues. Thereafter, the video takes a descending direction by giving some advice and suggestions (i.e., solutions) in the *Release*. They advise the viewers to stand up and stretch while in the office, then further encourage them to work out and be physically active by stating that physical activity is simple and can be done anytime and anywhere. The video uses walking as an example of simple but important physical activity that is free and easy to practice. The video concludes by providing concrete advice to practice 30 min of moderate-intensity physical activity 5 days per week, accompanied by images of a variety of suggested activities (e.g., basketball, hiking, etc.). We adjusted and used this video as follows:

- (1) **Condition 1** ‘Data Video with no solution’ (*baseline*): The video ends immediately after the peak, before any suggestions or solutions are provided (refer to video 1 in the supplementary material).
- (2) **Condition 2** ‘Data Video with broad/generalizable solutions’: This video provides a very general suggestion/solution after the peak, saying “physical activity is simple and can be done anywhere and anytime” (refer to video 2 in the supplementary material).
- (3) **Condition 3** ‘Data video with actionable/concrete solutions’: This video provides specific actionable solutions in addition to the general advice provided in video 2. Two concrete solution ideas were provided: walking (from the original video) and brisk walking (content and animation developed and added to the original video). To make the added advice useful and actionable, the video explains exactly what brisk walking is (e.g., “Brisk walking = 100 steps per minute or 17 steps per 10 seconds. Just walk fast enough so you can talk but not sing” (refer to video 3 in the supplementary material). Facts about brisk walking are taken from [73]⁹.

The film frames at the bottom of Figure 4 show a brief summary of the three videos.

4.3 Data Collection Instruments

Study 2 followed the same scales as Study 1 for measuring *personality traits* (Rammstedt and John’s [71] 10-item Big-Five Inventory) and *Persuasive potential* (Meschtscherjakov et al.’s [58] PPQ). Additionally, we measured:

Health Perception: We measured participants’ perception of their own physical activity using the statement “Generally speaking, I am being physically active” with a 7-point Likert scale: 1 indicating Strongly Disagree to 7, Strongly Agree.

Affective state: We adjusted the three questions in Study 1 to be focused on physical activity (see Table 2).

Actionability scale: Seven questions assessed the video content in terms of its utility or actionability (e.g., “Generally, the video provided concrete steps to address the problem.” on a 7-point Likert scale: 1 indicating Strongly Disagree to 7, Strongly Agree). These questions were created based on the ‘*Reduction*’ and ‘*Tunneling*’ principles from the Primary Task Support and ‘*Suggestions*’ from the Dialogue Support categories in Oinas-Kukkonen and Harjumaa’s [65] model for persuasive systems (see Appendix E for the actionability scale question items).

4.4 Overall Study Progression

First, participants answered demographic questions followed by the 10-item personality questions. Participants also answered a question regarding their physical activity level (“Generally speaking I am physically active”, on a 7-point Likert scale: 1 indicating Strongly Disagree to 7, Strongly Agree). Thereafter, participants started the video viewing section (see Appendix D). To capture the changes in participants’ affective reactions to the two solution videos, their

⁹To avoid any confounds, three HCI researchers ensured that the added content followed the same theme and animation style as the original video and also ensured smooth transitions between different scenes.

anxiety levels were assessed using the affective state questions (see Table 2) three times: 1) prior to the video viewing, 2) at the *Peak* point, and 3) after *Release* (i.e., Post video). For the control video, however, it was assessed twice: 1) prior to the video viewing (i.e., Pre-Video), 2) at the *Peak* point, because the video ended at its Peak point. Check Appendix D for a visual representation of the video viewing experience. After completing the video viewing, participants responded to:

Actionability Questions: measure perceived *utility or actionability* of the video (see Appendix E).

Video Content appraisal: assess perceived novelty, clarity, and usefulness of the DV content.

Health barriers: assesses whether it is possible for participants to follow the advice provided in the DV.

4.5 Results

The average completion time for Study 2 was roughly 15 minutes. Data fitting assumptions for each analysis were checked and non-parametric options were used when appropriate. Thus, medians (instead of means) are reported when applicable. Due to multiple comparisons, we used the Holm-Bonferroni method¹⁰. More detailed results, including null effects along with effect sizes, are reported in Appendix F.

4.5.1 Participants.

We recruited 119 participants¹¹ who varied in terms of gender, age, ethnicity, and education (see Table 3).

4.5.2 Data Quality Control.

The verifiable question used in Study 1 was asked again to filter out potentially unreliable responses. Applying a listwise deletion method for cases with wrong responses to the verifiable question left us with 109 valid cases out of 119; 8.4% of our data was considered invalid. We further filtered out responses when participants answered “yes” to the question “*I have health issues that prevent me from following the advice provided in the video*”. Eighty-two valid cases remained for our analyses ($n = 82$).

4.5.3 Data Preparation.

For each scale, we first checked inter-item reliability scores within the scale items (negatively coded items were reverse coded first) by investigating Cronbach’s *alphas*: 1) GPP ($\alpha = .84$); 2) Negative Affects ($\alpha = .95$); and 3) Actionability items ($\alpha = .89$). Since all scales encapsulate items that are sufficiently correlated (i.e., $\alpha \geq .70$ [63]), we created an aggregate index for each¹². Similarly, we created an index for each personality domain by aggregating its two items.

Negative Affect measures: In the study we measured participants’ negative affect two or three times depending on the condition (circular points on the line chart in Figure 4 represent times when negative affect was measured). Each time, we measured negative affect using the 3 anxiety questions in Table 2. After checking for

¹⁰The Holm-Bonferroni method was applied to balance both Type 1 and 2 errors. As the alternative to Bonferroni’s correction, where a consistent threshold is applied, Holm-Bonferroni’s method progressively adapts the threshold in which hypotheses are rejected based on the ranks of the mean scores of the comparisons.

¹¹Those people who participated in Study 1 were not recruited for Study 2.

¹²Whenever any of these four aspects is mentioned, they refer to the corresponding index.

inter-item reliability between the anxiety scale items, we created 3 indices:

- (1) **pre anxiety:** aggregate of participants’ anxiety level *before* watching any of the three video conditions.
- (2) **peak anxiety:** the aggregate of participants’ anxiety level *after watching the first segment of the video*, before solutions are presented (i.e., at the Peak).
- (3) **post anxiety:** the aggregate of participants’ anxiety level *after watching the second segment of the video* in which the solutions are presented (i.e., after the release in case of the solution video conditions).¹³

As the line graph in Figure 4 shows, participants in all three conditions exhibited elevated negative affect after viewing the first segment of the video which presented negative outcomes of not having enough physical activities. Note the change in anxiety levels cannot be attributed to the video condition yet because the videos were identical across all three conditions at this stage of the viewing (segment (a) in Figure 4). After exposing participants in the experimental conditions (i.e., the generalizable and actionable solution conditions) to the second segment of the video, in which the solutions are provided, their anxiety levels started to decline (post: Condition 2; $Mdn = 2.67$, Condition 3; $Mdn = 5.33$). As for participants in the control condition (i.e., the video with no solution), their anxiety level remained elevated as the video ended at its peak ($Mdn = 4.33$). For the analysis, we only considered the **Anxiety decrease/drop** per condition, which is the difference between median anxiety levels at the end of the video (post anxiety) and at the peak (peak anxiety) (see the red circles in Figure 4). This is the only measure that can be attributed to the condition effect.

4.5.4 Data Analysis and Findings.

Perceived Actionability: We checked if our video manipulation (i.e., adding two different types of solutions in the video) succeeded in making the videos more actionable. As Figure 5 shows, the trends are as anticipated. Next, a Kruskal-Wallis test examined if there is a significant difference between the three conditions in terms of actionability. We found a significant condition effect ($p = .004$). Pair-wise comparisons using Mann-Whitney U tests (details available in Appendix F) further found that participants in the experimental conditions perceived the videos as more actionable compared to participants in the control condition. Interestingly, there is no significant difference between the two experimental conditions (i.e., both of the experimental videos are significantly more actionable than the control video but are not different from each other)¹⁴. This indicates that, to improve perceived actionability, video designers can simply add a general solution to the video instead of detailed solutions. This is an interesting finding as adding concrete solutions usually requires more resources (e.g., detailed information search, more design effort, and more time added to the video).

Actionability and Usefulness: We examined whether actionability leads to information usefulness. As our nomological network suggested (see Appendix C), actionability accounted for the variation in usefulness, $F(1, 81) = 63.03$, $p < .001$, $\beta = .664$, $R^2_{adj} = .434$.

¹³For the video with no solution, post anxiety will be the same as peak anxiety as there is no second segment, so participants’ anxiety levels remain the same.

¹⁴Note, however, our sample size was smaller: for a better understanding of these null effects, please see Appendix F

Table 2: Physical Activity Negative Affect Question Items

Please read each statement carefully, and select the appropriate answer that best describes how you feel right now .
I feel worried about my health due to my low physical activity level.
I feel nervous about my physical activity level.
I feel anxious about my physical activity level.

A Likert scale (1: Not at all to 8: Extremely) is used.

Table 3: Participants' Demographic Information

Total Participants= 119	
Gender	Male (66%), Female (30%), Other (4%)
Age	18-34 (39%), 25-49 (40%), 50-64 (18%), over 65 (3%)
Ethnithity	White (77%), Asian, Black, Hispanic (16%), Other (5%), 2 did not respond
Education	High school or equivalent (17%), Bachelor's degree (61%), Graduate degree (19%), Other (3%)
Gotcha Question	Correct (109 participants) , Wrong (10 participants)
Health Issues	No (82), Yes (27)

Median Negative Affect Levels, GPP, and Usefulness per Video Condition




Figure 4: The yellow curve in the background is a rough representation of the common narrative arc and does not follow any scale. The line charts represent the fluctuation in participants' negative affect during the video viewing. Each line represents one video condition. The film frames at the bottom summarize the 3 video conditions and show where they end relative to each other and to the narrative arc as well as the affect fluctuation trends. (a) is identical across all three conditions, and it represents the first part of the videos in which the negative outcomes of not having enough physical activity are listed (E + I + P). The bar charts (right side) represent median GPP and median usefulness by video condition.




Figure 5: Medians of Perceived actionability per video condition. Error bars represent 95% CI.

Higher actionability predicted an increase in the usefulness of the information ('a' in Figure 6).

Usefulness: We also investigated the link between the video condition and perceived information usefulness. We entered the video condition as a predictor of information usefulness in a linear regression. The video condition predicted information usefulness ($F(1, 81) = 6.602, p = .012, \beta = .276, R^2_{adj} = .065$). The more actionable and detailed solutions provided in the video, the more useful it was considered. Further, a Kruskal-Wallis Test found that there was a significant difference in information usefulness among the three conditions ($p = 0.028$). Subsequent Mann-Whitney U Tests (details including effect sizes are available in Appendix F) found that, compared to the control video, the video with actionable solutions was perceived as significantly more useful ($p = 0.011$). No other significant effects were found. To improve perceived usefulness of information, thus, adding actionable solutions is essential (i.e., general solutions are insufficient).

Usefulness and GPP: Finally, we tested whether perceived information usefulness predicts general persuasive potential or GPP as found in Study 1. We entered usefulness as a predictor of GPP in a simple linear regression, and information usefulness did predict GPP ($F(1, 81) = 122.97, p < .001, \beta = .778, R^2_{adj} = .601$). Consistent with our findings in Study 1, information usefulness was a very strong predictor of GPP; it explained 60% of the variability in GPP ('b' in Figure 6).

Actionability, Anxiety, and GPP: To explore the effect of having solutions on easing participants' levels of negative affect, we conducted a linear regression with the video condition as a predictor of the level of drop in anxiety. The video condition predicted the drop in anxiety levels ($F(1, 81) = 63.03, p = .030, \beta = .240, R^2_{adj} = .046$). When the video provided a solution, the anxiety levels declined further, meaning that participants experienced more relief at the conclusion of the video. When no solution was provided,




Figure 6: Videos with solution are more actionable, actionable DVs predicted usefulness with $\beta = .664$. Usefulness predicted GPP with $\beta = .778$.

** : $p < .001$.

unlike the two solution conditions, participants' negative affect remained elevated at the end of the video. A pairwise comparison using the Mann-Whitney U test showed that there was no significant difference in the drop in anxiety levels between the two experimental conditions (generalizable and actionable solutions), please see Appendix F for more details. This finding informs us that incorporating *any* kind of solution in DVs can ease the anxiety that viewers develop from their exposure to the video. Next, we entered anxiety drop as a predictor of GPP in a linear regression model. Anxiety drop predicted GPP at marginal level¹⁵ ($F(1, 81) = 3.022, p = .086, \beta = .191, R^2 = .036$).




Figure 7: The video condition predicted a drop in negative affect (a); Video with detailed solutions induced the highest decrease in negative affect while video without solution did not decrease participants' negative affect. The drop in negative affect predicted GPP (b), marginally.

*: $P < .05$.

Solution and GPP: Lastly, we examined the effect of having a solution (generalizable solution or actionable solution) vs. no solution on overall GPP. For this analysis we split our data into only two conditions: 1) DV with no solution, 2) DV with solution (this included the general and actionable solution data). We entered the solution condition as a predictor of overall GPP in a linear regression model. Adding a solution did predict GPP ($\beta = .242, p = .028, R^2_{adj} = .047$). Videos with solutions (general or actionable) were seen as more persuasive than the video without a solution.

¹⁵We believe this marginal effect could have been due to a smaller N since we did find a small effect in this analysis. Note $R^2 = .036$ is considered to be a small effect according to Cohen [21]

Personal Aspects Influencing Negative Affect: To understand the dynamics underlying participants' personal tendencies to feel anxious or worried, we examined two factors:

Personality Traits and Negative Affect: With multiple linear regressions using a stepwise method, we explored personality traits as predictors of overall negative affect index (mean of pre, peak, and post anxiety). We were able to replicate the findings from Study 1: only Conscientiousness ($\beta = -.388, p < .001$) and Neuroticism ($\beta = .358, p = .001$) predicted overall negative affect $F(2, 80) = 68.64, p < .001, R^2_{adj} = .401$.

Health Perception and Negative Affect: We also looked at the relationship between participants' perceived physical activity levels and the anxiety levels they reported after watching the "threatening" part of the video (i.e., peak anxiety). A linear regression analysis found that participants' health perception predicted negative affect with a negative *beta* ($\beta = -.316, p = .004, R^2_{adj} = .088$). When participants perceived themselves as less physically active they felt more anxious and worried after watching the video.

These findings tell us that highly neurotic people are more susceptible to fear-inducing messages while people who score high in conscientiousness are more resilient to such messages. Also, when participants perceived themselves as less physically active (i.e., presumably the targets of such DVs), they tend to become more anxious when presented with threatening (or loss-framed) messages related to physical inactivity. Thus, designing different versions of DVs based on the viewers' personality traits, specifically on neuroticism and conscientiousness, may motivate viewers more efficiently.

5 DISCUSSION

To get a better understanding of the persuasive potential of Data Videos targeting common health issues, this research focused on the exploration of viewers' affect, personality, the type of provided information, and the structure of Data Videos. The first study asked three questions; how is viewers' affect linked to their willingness to alter their health-related behaviours?; how are these affective responses linked to the viewers' personality traits?; and what are the underlying attributes in DVs that contribute to their persuasive potential? With the findings of Study 1, Study 2 then focused on providing guidelines for video designers by exploring how we can adjust participants' affect along with the narrative structure to improve the overall impact of DVs by altering the *types of information* provided. In this section, we first discuss how our findings are related to and explained by, previous work. We also present a set of initial guidelines which designers of DVs can follow.

We observed that DVs lacking clear solutions to their loss-framed messages kept viewers' negative affect higher after their video viewing. Then, these heightened levels of negative affect predicted a lower viewer willingness to change their behaviour. Moreover, the viewers' levels of negative affect were linked to their own personality traits. While our results cannot tell us *how exactly* we should design trait-specific DVs, they flag the potential risk of including heavily loss-framed messages. While loss-framed messages are very commonly included in DVs, and the relationship between fear and persuasion has been validated in numerous research [25, 53], in the

context of DVs, such fear/anxiety-inducing messages could potentially backfire; The message could be perceived as less convincing if viewers' negative affect were left unresolved by the end of the video. Interestingly, our results are consistent with earlier psychological investigations of smoking. For example, one study revealed that fear was effective in *motivating* behaviour change, but only when the message informed individuals how to succeed in smoking cessation efforts; loss-framed message with solutions[52].

This research provides design guidelines (GLs) to improve the persuasive potential of DVs targeting health issues in two directions: 1) the content and narrative structure of the videos; and 2) personality-tailored guidelines.

5.1 Data Video Content: Related to RQ1 & 3

To increase the persuasive potential of health-related DVs that normally follow a loss-framed messaging style (i.e., messages that show the consequences of failing to take certain actions), designers should make sure to follow a '*well-balanced*' narrative structure: Presenting the problem and all the threats associated with it, should be followed by solution(s) or suggestions(s) to balance the level of negative affect viewers experience throughout the video viewing, and to leave the viewers emotionally resolved by the end of the DV (Study 1). Our studies proved that including solutions in health-related DVs makes the viewers perceive the videos as more actionable and useful, eases viewers' anxiety levels, and makes the videos more persuasive. Importantly, we observed trends around solution type: while presenting simple general solutions was effective in reducing anxiety levels and improving the actionability of DVs, detailed actionable solutions that provide very specific advice consistently yielded the best outcomes (Study 2).

GL1: *DVs targeting health issues should include a solution to the presented problem. This should leave the viewers emotionally resolved by nicely balancing the levels of negative affect they experienced throughout the video viewing.*

GL2: *Adding detailed and specific solutions to DVs is clearly an effective way to make them useful and less stressful. However, if the video length is an issue, while not ideal, adding a simple solution would also improve the power of DVs.*

5.2 Personalization of DVs: Related to RQ1 & 2

As for our contributions in terms of personalization of DVs, our findings confirm that when watching DVs which target a health issue, people react differently to affect-provoking messages due to their underlying personality traits. Highly neurotic individuals experience higher levels of negative affect when exposed to threatening messages, while highly conscientious individuals have a higher tolerance for receiving such messages. Neurotic people are also less likely to be persuaded via watching DVs. Furthermore, people who perceive themselves as more vulnerable to the health issue discussed in the video (i.e., possibly the target of DVs) are more sensitive to the threatening messages.

GL3: *Designers should be extra cautious when targeting individuals with neurotic tendencies in terms of the amount and/or levels of threat contained in their messages or else their DVs may backfire.*

GL4: *If designers are targeting individuals who feel vulnerable to the health issue, they should limit the amount and/or levels of threat in the messages.*

Two studies confirmed that DVs have the potential to improve viewers' behavioural change intentions. Given their popularity, DVs should be carefully designed in order to use them to their full potential. This research provides some guidelines that should help designers of DVs targeting health issues.

6 LIMITATIONS AND FUTURE WORK

We were not able to conduct in-person studies due to the COVID-19 pandemic; instead, we conducted online studies, which limited the controllability of our study. This remote setup also meant we had to rely on participants' self-reporting of their affective responses. Future studies with physiological measurements to assess affective responses will be useful to validate our findings. Furthermore, the studies were limited to negative affective responses focusing on anxiety, which may have been affected by the pandemic as participants might have been experiencing higher levels of general anxiety than usual. Additionally, the investigation of positive affect (e.g., excitement, hope) will also improve our model further. This could lead us to guidelines on how to design effective data videos using positive framing, rather than by inducing fear, especially for those who score high on neuroticism. Our studies shed light on the importance of some parts of perceived usefulness of the content in Data Videos (i.e., actionability). Future studies should investigate other dimensions of perceived usefulness in the content of DVs. With respect to personality, we followed the Big-5 personality traits. Future research might also look to other traits, such as authoritarianism [3], to understand how those aspects interact with Data Video effectiveness. We acknowledge that our results could be limited to the common health topics covered in the videos and future studies should also explore other less common health issues for which the DVs can provide their viewers with new information. Additionally, our studies had rather smaller sample sizes, and this could have led us to reject the null effects. To compensate for this issue, we provided effect sizes (which are independent from sample size issues [84]) and the descriptions of null effects in the appendices to provide a clearer picture of our results. While some effects we found in Study 1 were replicated in Study 2, another study with a greater sample size will be important. Finally, our studies looked at behaviour change on the perceptual level as opposed to actual behaviour change. A longitudinal study to explore people's actual behavioural change will be fruitful.

7 CONCLUSION

Data Videos (DV) which convey rich and complex information in a short period of time, are a relatively new form of data representation that has become prominent on social media. DVs are gaining in popularity and their viewing rates are skyrocketing [12], hence they constitute a promising medium for data representation which is worth exploring. The design and creation of effective DVs require aesthetic design skills and reliable information sources. In order to move towards personalized DVs, designers also require an empirical understanding of how viewers' personalities and affective states interact with the video content and framing. Our research

contributes empirical evidence about these interactions which can help video designers create DVs that are more personalized and effective. We conducted two empirical studies online, which involved participants watching health-related DVs and responding to a variety of survey instruments before, after, and in between video segments. We explored viewers' affective reactions in response to loss-framed messages in DVs and how those affective reactions were linked to the viewers' unique personalities as well as viewers' potential to consider changing their related health behaviours. Our two studies generated guidelines that DV designers can follow to create more effective, personalized health-related DVs. Our findings may also contribute to the development of novel platforms for personalized self-help and behavioural change, such as interactive DVs.

ACKNOWLEDGMENTS

This study was supported by NSERC. We also thank Sepideh Saffari for her help in the creation of the illustrations used in the teaser image (Figure 1).

REFERENCES

- [1] 2018. Physical activity. <https://www.who.int/news-room/fact-sheets/detail/physical-activity>. <https://www.who.int/news-room/fact-sheets/detail/physical-activity>
- [2] David A Aaker and Donald E Bruzzone. 1981. Viewer perceptions of prime-time television advertising. *Journal of Advertising research* (1981).
- [3] Theodor Adorno, Else Frenkel-Brenswik, Daniel J Levinson, and R Nevitt Sanford. 2019. *The authoritarian personality*. Verso Books.
- [4] Haldun Akoglu. 2018. User's guide to correlation coefficients. *Turkish journal of emergency medicine* 18, 3 (2018), 91–93.
- [5] Fereshteh Amini, Nathalie Henry Riche, Bongshin Lee, Christophe Hurter, and Pourang Irani. 2015. Understanding data videos: Looking at narrative visualization through the cinematography lens. In *Proceedings of the 33rd Annual ACM Conference on Human Factors in Computing Systems*. 1459–1468.
- [6] Fereshteh Amini, Nathalie Henry Riche, Bongshin Lee, Jason Leboe-McGowan, and Pourang Irani. 2018. Hooked on Data Videos: Assessing the Effect of Animation and Pictographs on Viewer Engagement. In *Proceedings of the 2018 International Conference on Advanced Visual Interfaces (Castiglione della Pescaia, Grosseto, Italy) (AVI '18)*. ACM, New York, NY, USA, Article 21, 9 pages. <https://doi.org/10.1145/3206505.3206552>
- [7] Fereshteh Amini, Nathalie Henry Riche, Bongshin Lee, Andres Monroy-Hernandez, and Pourang Irani. 2016. Authoring data-driven videos with dataclips. *IEEE transactions on visualization and computer graphics* 23, 1 (2016), 501–510.
- [8] Sonia M Arteaga, Mo Kudeki, and Adrienne Woodworth. 2009. Combating obesity trends in teenagers through persuasive mobile technology. *ACM SIGACCESS Accessibility and Computing* 94 (2009), 17–25.
- [9] Chris Baber, Dan Andrews, Tom Duffy, and Richard McMaster. 2011. Sensemaking as narrative: Visualization for collaboration. *VAW2011, University London College* (2011), 7–8.
- [10] Shaowen Bardzell, Jeffrey Bardzell, and Tyler Pace. 2009. Understanding affective interaction: Emotion, engagement, and internet videos. In *2009 3rd International Conference on Affective Computing and Intelligent Interaction and Workshops*. IEEE, 1–8.
- [11] John A Bargh, Peter M Gollwitzer, Annette Lee-Chai, Kimberly Barndollar, and Roman Trötschel. 2001. The automated will: nonconscious activation and pursuit of behavioral goals. *Journal of personality and social psychology* 81, 6 (2001), 1014.
- [12] Corey H Basch, Grace C Hillyer, Emily A Zagnit, and Charles E Basch. 2020. YouTube coverage of COVID-19 vaccine development: implications for awareness and uptake. *Human vaccines & immunotherapeutics* 16, 11 (2020), 2582–2585.
- [13] Trevor Bopp, Joshua D Vadeboncoer, Michael Stelfelson, and Melissa Weinsz. 2019. Moving beyond the gym: A content analysis of YouTube as an information resource for physical literacy. *International Journal of Environmental Research and Public Health* 16, 18 (2019), 3335.
- [14] Ron Borland, Hua-Hie Yong, Nick Wilson, Geoffrey T Fong, David Hammond, K Michael Cummings, Warwick Hosking, and Ann McNeill. 2009. How reactions to cigarette packet health warnings influence quitting: Findings from the ITC Four-Country survey. *Addiction* 104, 4 (2009), 669–675.
- [15] Kathryn A Braun. 1999. Postexperience advertising effects on consumer memory. *Journal of consumer research* 25, 4 (1999), 319–334.

- [16] Marian Chapman Burke and Julie A Edell. 1989. The impact of feelings on ad-based affect and cognition. *Journal of marketing research* 26, 1 (1989), 69–83.
- [17] John T Cacioppo and Richard E Petty. 1984. The elaboration likelihood model of persuasion. *ACR North American Advances* (1984).
- [18] Ruochen Cao, Subrata Dey, Andrew Cunningham, James Walsh, Ross T Smith, Joanne E Zucco, and Bruce H Thomas. 2019. Examining the use of narrative constructs in data videos. *Visual Informatics* (2019).
- [19] Chun-Tuan Chang and Yu-Kang Lee. 2011. The 'I' of the beholder: How gender differences and self-referencing influence charity advertising. *International Journal of Advertising* 30, 3 (2011), 447–478.
- [20] Eun Kyoung Choe, Yumiko Sakamoto, Yanis Fatmi, Bongshin Lee, Christophe Hurter, Ashkan Haghshenas, and Pourang Irani. 2019. Persuasive Data Videos: Investigating Persuasive Self-Tracking Feedback with Augmented Data Videos. In *AMIA Annual Symposium Proceedings*, Vol. 2019. American Medical Informatics Association, 295.
- [21] Jacob Cohen. 1988. Statistical power analysis for the behavioral sciences. Hillsdale, NJ: Lawrence Erlbaum Associates. Inc. Publishers (1988).
- [22] Jonathan Cohen. 2001. Defining identification: A theoretical look at the identification of audiences with media characters. *Mass communication & society* 4, 3 (2001), 245–264.
- [23] Neil Cohn. 2013. Visual narrative structure. *Cognitive science* 37, 3 (2013), 413–452.
- [24] Annelise L Dawson, Ashley A Hamstra, Laura Sturgess Huff, Ryan G Gamble, William Howe, Ilima Kane, and Robert P Dellavalle. 2011. Online videos to promote sun safety: results of a contest. *Dermatology Reports* 3, 1 (2011).
- [25] James Price Dillard and Jason W Anderson. 2004. The role of fear in persuasion. *Psychology & Marketing* 21, 11 (2004), 909–926.
- [26] Amy M Do, Alexander V Rupert, and George Wolford. 2008. Evaluations of pleasurable experiences: The peak-end rule. *Psychonomic Bulletin & Review* 15, 1 (2008), 96–98.
- [27] Filip Drozd, Tuomas Lehto, and Harri Oinas-Kukkonen. 2012. Exploring perceived persuasiveness of a behavior change support system: a structural model. In *International Conference on Persuasive Technology*. Springer, 157–168.
- [28] Ian Duncan, Lee Yarwood-Ross, and Carol Haigh. 2013. YouTube as a source of clinical skills education. *Nurse education today* 33, 12 (2013), 1576–1580.
- [29] Hans J Eysenck. 1992. Four ways five factors are not basic. *Personality and individual differences* 13, 6 (1992), 667–673.
- [30] World Health Organization Regional Office for the Western Pacific. 2016. Move for Health #2. <https://www.youtube.com/watch?v=OxeFwv4AeKM&t=47s>
- [31] Marilyn Frenn, Shelly Malin, Roger L Brown, Yvonne Greer, Jaime Fox, Jennifer Greer, and Sarah Smyczek. 2005. Changing the tide: an Internet/video exercise and low-fat diet intervention with middle-school students. *Applied nursing research* 18, 1 (2005), 13–21.
- [32] Lena Frischlich, Diana Rieger, Anna Morten, and Gary Bente. 2018. The power of a good story: Narrative persuasion in extremist propaganda and videos against violent extremism. *International Journal of Conflict and Violence (IJCV)* 12 (2018), a644–a644.
- [33] Elia Gabarron, Luis Fernandez-Luque, Manuel Armayones, and Annie YS Lau. 2013. Identifying measures used for assessing quality of YouTube videos with patient health information: a review of current literature. *Interactive journal of medical research* 2, 1 (2013), e6.
- [34] Kristel M Gallagher and John A Updegraff. 2012. Health message framing effects on attitudes, intentions, and behavior: a meta-analytic review. *Annals of behavioral medicine* 43, 1 (2012), 101–116.
- [35] Selin Gaş, Özge Ö Zincer, and Aylin P Bozkurt. 2019. Are YouTube videos useful for patients interested in botulinum toxin for bruxism? *Journal of Oral and Maxillofacial Surgery* 77, 9 (2019), 1776–1783.
- [36] Alan S Gerber, Gregory A Huber, David Doherty, Conor M Dowling, and Costas Panagopoulos. 2013. Big five personality traits and responses to persuasive appeals: Results from voter turnout experiments. *Political Behavior* 35, 4 (2013), 687–728.
- [37] Nahum Gershon and Ward Page. 2001. What storytelling can do for information visualization. *Commun. ACM* 44, 8 (2001), 31–37.
- [38] Marvin E Goldberg and Gerald J Gorn. 1987. Happy and sad TV programs: How they affect reactions to commercials. *Journal of consumer research* 14, 3 (1987), 387–403.
- [39] Samuel D Gosling, Peter J Rentfrow, and William B Swann Jr. 2003. A very brief measure of the Big-Five personality domains. *Journal of Research in personality* 37, 6 (2003), 504–528.
- [40] Melanie C Green and Timothy C Brock. 2000. The role of transportation in the persuasiveness of public narratives. *Journal of personality and social psychology* 79, 5 (2000), 701.
- [41] Philip J Guo, Juho Kim, and Rob Rubin. 2014. How video production affects student engagement: An empirical study of MOOC videos. In *Proceedings of the first ACM conference on Learning@ scale conference*. 41–50.
- [42] Sajaneh Halko and Julie A Kientz. 2010. Personality and persuasive technology: an exploratory study on health-promoting mobile applications. In *International conference on persuasive technology*. Springer, 150–161.
- [43] Leslie J Hinyard and Matthew W Kreuter. 2007. Using narrative communication as a tool for health behavior change: a conceptual, theoretical, and empirical overview. *Health Education & Behavior* 34, 5 (2007), 777–792.
- [44] Jacob B Hirsh, Sonia K Kang, and Galen V Bodenhausen. 2012. Personalized persuasion: Tailoring persuasive appeals to recipients' personality traits. *Psychological science* 23, 6 (2012), 578–581.
- [45] Morris B Holbrook and Rajeev Batra. 1987. Assessing the role of emotions as mediators of consumer responses to advertising. *Journal of consumer research* 14, 3 (1987), 404–420.
- [46] Maurits Kaptein, Panos Markopoulos, Boris De Ruyter, and Emile Aarts. 2015. Personalizing persuasive technologies: Explicit and implicit personalization using persuasion profiles. *International Journal of Human-Computer Studies* 77 (2015), 38–51.
- [47] Maurits Clemens Kaptein. 2012. Personalized persuasion in ambient intelligence. (2012).
- [48] Robert Kosara and Jock Mackinlay. 2013. Storytelling: The next step for visualization. *Computer* 46, 5 (2013), 44–50.
- [49] Michael Kubovy, Daniel Kahneman, Ed Diener, N Schwarz, et al. 1999. Well-being: The foundations of hedonic psychology. (1999).
- [50] Celine Latulipe, Erin A Carroll, and Danielle Lottridge. 2011. Love, hate, arousal and engagement: exploring audience responses to performing arts. In *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*. 1845–1854.
- [51] Loraine Lau-Gesk and Joan Meyers-Levy. 2009. Emotional persuasion: When the valence versus the resource demands of emotions influence consumers' attitudes. *Journal of Consumer Research* 36, 4 (2009), 585–599.
- [52] Howard Leventhal, Jean C Watts, and Franca Pagano. 1967. Effects of fear and instructions on how to cope with danger. *Journal of personality and social psychology* 6, 3 (1967), 313.
- [53] Ioni Lewis, Barry Watson, and Katherine M White. 2013. Extending the explanatory utility of the EPPM beyond fear-based persuasion. *Health communication* 28, 1 (2013), 84–98.
- [54] Elizabeth J Marsh. 2006. When does generation enhance memory for location? *Journal of Experimental Psychology: Learning, Memory, and Cognition* 32, 5 (2006), 1216.
- [55] Theresa M Marteau and Hilary Bekker. 1992. The development of a six-item short-form of the state scale of the Spielberger State–Trait Anxiety Inventory (STAI). *British journal of clinical Psychology* 31, 3 (1992), 301–306.
- [56] Sandra C Matz, Michal Kosinski, Gideon Nave, and David J Stillwell. 2017. Psychological targeting as an effective approach to digital mass persuasion. *Proceedings of the national academy of sciences* 114, 48 (2017), 12714–12719.
- [57] Robert R McCrae and Oliver P John. 1992. An introduction to the five-factor model and its applications. *Journal of personality* 60, 2 (1992), 175–215.
- [58] Alexander Meschtscherjakov, Magdalena Gärtner, Alexander Mirnig, Christina Rödel, and Manfred Tscheligi. 2016. The persuasive potential questionnaire (ppq): Challenges, drawbacks, and lessons learned. In *International Conference on Persuasive Technology*. Springer, 162–175.
- [59] Emily Moyer-Gusé. 2008. Toward a theory of entertainment persuasion: Explaining the persuasive effects of entertainment-education messages. *Communication theory* 18, 3 (2008), 407–425.
- [60] Elizabeth L Murnane, Xin Jiang, Anna Kong, Michelle Park, Weili Shi, Connor Soohoo, Luke Vink, Iris Xia, Xin Yu, John Yang-Sammataro, et al. 2020. Designing Ambient Narrative-Based Interfaces to Reflect and Motivate Physical Activity. In *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems*. 1–14.
- [61] Sheila T Murphy, Lauren B Frank, Meghan B Moran, and Paula Patnoe-Woodley. 2011. Involved, transported, or emotional? Exploring the determinants of change in knowledge, attitudes, and behavior in entertainment-education. *Journal of communication* 61, 3 (2011), 407–431.
- [62] John P Murry Jr and Peter A Dacin. 1996. Cognitive moderators of negative-emotion effects: Implications for understanding media context. *Journal of Consumer Research* 22, 4 (1996), 439–447.
- [63] Jum C Nunnally and IH Bernstein. 1978. Psychometric Theory McGraw-Hill New York. *The role of university in the development of entrepreneurial vocations: a Spanish study* (1978).
- [64] Institute of Medicine (US). Committee on Health, Practice, and Policy. 2001. *Health and behavior: The interplay of biological, behavioral, and societal influences*. National Academies Press.
- [65] Harri Oinas-Kukkonen and Marja Harjuma. 2009. Persuasive systems design: Key issues, process model, and system features. *Communications of the Association for Information Systems* 24, 1 (2009), 28.
- [66] Rita Orji, Lennart E Nacke, and Chrysanne Di Marco. 2017. Towards personality-driven persuasive health games and gamified systems. In *Proceedings of the 2017 CHI Conference on Human Factors in Computing Systems*. 1015–1027.
- [67] Gjalit-Jorn Ygram Peters, Robert AC Ruiter, and Gerjo Kok. 2013. Threatening communication: a critical re-analysis and a revised meta-analytic test of fear appeal theory. *Health psychology review* 7, sup1 (2013), S8–S31.
- [68] Janey C Peterson, Mary E Charlson, Zachary Hoffman, Martin T Wells, Shing-Chiu Wong, James P Hollenberg, Jared B Jobe, Kathryn A Boschert, Alice M Isen,

- and John P Allegrante. 2012. A randomized controlled trial of positive-affect induction to promote physical activity after percutaneous coronary intervention. *Archives of internal medicine* 172, 4 (2012), 329–336.
- [69] Richard E Petty, Joseph R Priester, and Duane T Wegener. 1994. Cognitive processes in attitude change. *Handbook of social cognition* 2 (1994), 69–142.
- [70] Tom Pyszczynski, Jeff Greenberg, and Sheldon Solomon. 1999. A dual-process model of defense against conscious and unconscious death-related thoughts: an extension of terror management theory. *Psychological review* 106, 4 (1999), 835.
- [71] Beatrice Rammstedt and Oliver P John. 2007. Measuring personality in one minute or less: A 10-item short version of the Big Five Inventory in English and German. *Journal of research in Personality* 41, 1 (2007), 203–212.
- [72] Donald A Redelmeier and Daniel Kahneman. 1996. Patients' memories of painful medical treatments: Real-time and retrospective evaluations of two minimally invasive procedures. *pain* 66, 1 (1996), 3–8.
- [73] Gretchen Reynolds. 2018. Walk briskly for your Health. about 100 steps a minute. <https://www.nytimes.com/2018/06/27/well/walk-health-exercise-steps.html>
- [74] Alexander J Rothman and Peter Salovey. 1997. Shaping perceptions to motivate healthy behavior: the role of message framing. *Psychological bulletin* 121, 1 (1997), 3.
- [75] Margaret Sampson, Jordi Cumber, Claudia Li, Catherine M Pound, Ann Fuller, and Denise Harrison. 2013. A systematic review of methods for studying consumer health YouTube videos, with implications for systematic reviews. *PeerJ* 1 (2013), e147.
- [76] Paul Sărbescu and Alexandru Boncu. 2018. The resilient, the restraint and the restless: Personality types based on the Alternative Five-Factor Model. *Personality and Individual Differences* 134 (2018), 81–87.
- [77] Edward Segel and Jeffrey Heer. 2010. Narrative visualization: Telling stories with data. *IEEE transactions on visualization and computer graphics* 16, 6 (2010), 1139–1148.
- [78] Avi Shankar, Richard Elliott, and Christina Goulding. 2001. Understanding consumption: Contributions from a narrative perspective. *Journal of marketing Management* 17, 3-4 (2001), 429–453.
- [79] Matthew A Shapiro and Han Woo Park. 2015. More than entertainment: YouTube and public responses to the science of global warming and climate change. *Social Science Information* 54, 1 (2015), 115–145.
- [80] Michael D Slater. 1997. Persuasion processes across receiver goals and message genres. *Communication Theory* 7, 2 (1997), 125–148.
- [81] Michael D Slater and Donna Rouner. 2002. Entertainment—education and elaboration likelihood: Understanding the processing of narrative persuasion. *Communication theory* 12, 2 (2002), 173–191.
- [82] Joanne R Smith, Michael A Hogg, Robin Martin, and Deborah J Terry. 2007. Uncertainty and the influence of group norms in the attitude-behaviour relationship. *British Journal of Social Psychology* 46, 4 (2007), 769–792.
- [83] Mohammad Soleymani, Sander Koelstra, Ioannis Patras, and Thierry Pun. 2011. Continuous emotion detection in response to music videos. In *Face and Gesture 2011*. IEEE, 803–808.
- [84] Gail M Sullivan and Richard Feinn. 2012. Using effect size—or why the P value is not enough. *Journal of graduate medical education* 4, 3 (2012), 279–282.
- [85] Sean Tackett, Kyle Slinn, Tanner Marshall, Shiv Gaglani, Vincent Waldman, and Rishi Desai. 2018. Medical education videos for the world: an analysis of viewing patterns for a YouTube channel. *Academic medicine* 93, 8 (2018), 1150–1156.
- [86] Amos Tversky and Daniel Kahneman. 1985. The framing of decisions and the psychology of choice. In *Behavioral decision making*. Springer, 25–41.
- [87] Tom Van Laer, Ko De Ruyter, Luca M Visconti, and Martin Wetzels. 2014. The extended transportation-imagery model: A meta-analysis of the antecedents and consequences of consumers' narrative transportation. *Journal of Consumer research* 40, 5 (2014), 797–817.
- [88] Helen J Wall, Claire C Campbell, Linda K Kaye, Andy Levy, and Navjot Bhullar. 2019. Personality profiles and persuasion: An exploratory study investigating the role of the Big-5, Type D personality and the Dark Triad on susceptibility to persuasion. *Personality and Individual Differences* 139 (2019), 69–76.
- [89] Wibke Weber. 2020. Exploring narrativity in data visualization in journalism. *Data visualization in society* (2020), 295–311.
- [90] Wibke Weber, Martin Engebretsen, and Helen Kennedy. 2018. Data stories: Rethinking journalistic storytelling in the context of data journalism. *Studies in Communication Sciences* 2018, 1 (2018), 191–206.
- [91] Jessica Wortman and Dustin Wood. 2011. The personality traits of liked people. *Journal of Research in Personality* 45, 6 (2011), 519–528.
- [92] Li-fang Zhang and Jiafen Huang. 2001. Thinking styles and the five-factor model of personality. *European Journal of Personality* 15, 6 (2001), 465–476.

A LIST OF THE NINE DATA VIDEOS USED IN THE STUDY WITH THEIR RELEVANT LINKS AND SOME ADDITIONAL INFORMATION

Table 4: List of the Data Videos used in the study, by topic

Title	URL	Source	Length	# Views	# Data Representations	Audio Narration
Physical Activity						
Morning Exercise = Mental Health	https://www.youtube.com/watch?v=3ZBXldCxZEA	Kram Gallery	2:12	373,174	1	yes
The Importance of Intensity in Physical Activity	https://www.youtube.com/watch?v=OMn8Tq5Eyao	DocMikeEvans	3:20	97,728	4	yes
What Happens To Your Body When You Stop Exercising	https://www.youtube.com/watch?v=hQz_V9Dr8IU	Tech Insider	2:08	6,339,611	8	no
Diet						
Sugar is Killing Us	https://www.youtube.com/watch?v=Yda8RtOcVFU	Sikuvideo	3:23	1,228,701	3	yes
What would happen if you didn't drink water?	https://www.youtube.com/watch?v=9iMGFqMmUFs	TED-ED	4:36	9,027,233	11	yes
Why eating fat won't make you gain weight	https://www.youtube.com/watch?v=47O619aXIt0	Tech Insider	4:43	346,114	8	yes
Sleep						
Understanding the Importance of Sleep	https://www.youtube.com/watch?v=NwJ_Xwr2irI	Tylenolofficial	1:25	28,524	3	yes
The benefits of a good night's sleep	https://www.youtube.com/watch?v=gedoSfZvBgE	Ted-Ed	5:30	3,445,468	3	yes
What causes insomnia?	https://www.youtube.com/watch?v=j5Sl8LyI7k8	Ted-Ed	4:47	1,930,075	6	yes

B LIST OF THE DATA VIDEOS USED WITH SUMMARY DESCRIPTION AND SOME VIDEO ATTRIBUTES










Video	Summary	Negative Consequences	Concrete Solution	# of Infographics
Physical Activity				
 <p>PA1: Morning Exercise = Mental Health 2:13</p>	The video is focusing on the importance of starting the day with physical activity. Benefits of exercising in the morning as: - mental capacity for handling stress triples, - the brain produces chemicals (BNF and Endorphins) that clears the mind, helps in making positive decisions, minimize the discomfort and the feeling of pain and promotes the feeling of pleasure - helps with a faster metabolism, more energy and sense of accomplishment.	✗	✓	1
 <p>PA2: The Importance of Intensity in Physical Activity 3:28</p>	- Narrator is Dr. Mike Evans (which adds credibility and authority to the video). - Video is about the importance of intensity in physical activity and they define it as "stuff that gets your heart pumping" - They state that we need to perform 150 minutes of Moderate to Vigorous Physical Activity weekly. - Exercise reduces heart disease and cancer, and improves mental health and the overall quality of our lives. - Research has shown that higher the intensity, the better the results. - Interval training is good to improve performance in a short amount of time. Tips to know the intensity of the activity: 1. low intensity activity: if you are not sweating or not experiencing breath reate change. 2. Moderate activity: you feel out of breath but can still hold a conversation 3. Vigorous activity: you can't talk"	✗	✓	4
 <p>PA3: What Happens to Your Body When You Stop Exercising 2:09</p>	- The video is talking about changes and risks to the body over time as we stop exercising. - The video is reporting changes that can happen over weeks to months. - The video is tracking the increase in the time needed to finish a 5k run (starting at 20 min when physically active, increase of 1 min after 2-3 weeks and 3 minutes longer after 5-7 weeks. - The video is also showing the decrease in the level of VO2 max that is responsible for delivering oxygen to generate energy for muscles as the no exercise period increases. As VO2 max decreases, muscle cells become smaller and the fat cells are become bigger.	✓	✗	8
Diet				
 <p>Diet1: Sugar is Killing Us 3:31</p>	- The video is talking about the dangers of sugar. - Scientists are saying sugar is toxic. It turned out that sugar is not only not healthy but it is dangerous. - High fructose in sugar can harm the body in various ways and causes lots of health issues. - However it is not easy to cut down on sweets as high fructose corn syrup are now added to foods that you never expect. - A recent study found the 80% of food in America contain added sugar. - At the end, the video is suggesting that the solution is to educate consumers not to buy products with added sugar which will affect the sales of such products.	✓	✗	3
 <p>WHAT WOULD HAPPEN IF YOU DIDN'T DRINK WATER? 4:52</p>	- The video is about the role of water in the human body and how much do we need to drink to stay healthy. - Water cushion and lubricate joints, regulate temperature and nourish the brain and spinal cord. - Each day our bodies lose 2-3 liters of water through sweat, urine and bowel movements and even from breathing. We need to drink water to compensate for this loss. - The amount of daily water intake we need ranges between 2.5-3.7 L for men and 2-2.7 L for women depending on the weight and environment. - Benefits of being well hydrated: lower the chance of stroke, manage diabetes, reduce the risk of certain types of cancer	✓	✓	11
 <p>WHY FAT ISN'T BAD TECH INSIDER 4:44</p>	- This video is focused on the benefits of good fats and how they are not associated with weight gain while carbs are the real issue. - Research shows that low fat diets do not aid in weight loss or reducing the risk of disease compared to high-fat diet and the refined carbs might be the real issue. - Good fats are very important for the body to function properly. - Monosaturated fats are found in olive oil and avocados and they reduce inflammations and levels of LDL or bad cholesterol in the blood. - Polyunsaturated fats are found in fish are very useful, Omega-3 fatty acids decrease blood pressure and increase DHL, or good Cholesterol and protects against heart disease. - Studies showed that full fat dairy is healthier than reduced fat as full fat dairy is found to be associated with a lower risk of type 2 diabetes	✓	✓	8
Sleep				
 <p>Sleep1: Understanding the Importance of Sleep 1:26</p>	This video is short and to the point. It explains the importance of getting good sleep, how much sleep do we need and the dangers of not getting enough sleep. Almost half of the video is dedicated to listing annoying and worrying threats of inadequate sleep (e.g. you will look older, you will get fat, you will develop cardiovascular disease, etc.)	✓	✗	3
 <p>SLEEP TO REMEMBER REMEMBER TO SLEEP 5:45</p>	- The video is talking about the importance of sleep for memory consolidation. - It starts by putting the viewers in an imaginary situation where they have a test in 8 hours followed by a piano recital and you are not feeling ready yet. You can drink a cup of coffee and spend some time studying and practicing but the narrator assures that it is better to put everything away and sleep. - The video contains a lot of scientific informations and terms and refers to several scientific studies related to the brain and the process of memory consolidation.	✓	✓	3
 <p>WHAT CAUSES INSOMNIA? 5:12</p>	- The video focuses on insomnia, its symptoms, reasons and provides some solutions. - unlike healthy sleepers, during their sleep, insomniac experience high metabolism which exhausts their bodies and brain making them wake in a state exhaustion, stress and confusion. - Insomnia's chemical mechanisms are similar to anxiety attacks found in people experiencing depression and anxiety and accordingly suffering from insomnia increases the risk of suffering from depression and anxiety.	✓	✓	6

Figure 8: List of all videos with summary description and some video attributes.

C NOMOLOGICAL NETWORK ON INFORMATION USEFULNESS IN DATA VIDEOS




Figure 9: Nomological network of the dimensions of Information usefulness in data videos

D VIDEO VIEWING EXPERIENCE




Figure 10: Participants were randomly assigned to one of the three video conditions:

1. In the *No Solution* condition, participants watched one video segment that covered the dangers of not being physically active then answered the Affective state questions.
2. In the *General Solution* Condition, participants watched 2 video segments; the first segment covered the dangers of not being physically (identical to condition 1), then they answered the affective state question, finally they watched the second video segment that introduced a general solution.
3. In the *Actionable Solution* Condition, participants watched 2 video segments; the first segment covered the dangers of not being physically (identical to condition 1 and 2), then they answered the affective state question, finally they watched the second video segment that introduced an actionable solution.

E ACTIONABILITY SCALE

Table 5: Items to measure the videos' actionability

Please report you thoughts about the video.

Generally, the video provided solution(s) to the discussed issue.
Generally, the video provided concrete steps to address the problem.
Generally, the video provided easy to understand solutions
Generally, people would be capable of implementing the proposed solutions.
Generally, people would have the required resources (time, space, etc.) to carry out the suggestions in the video.
Generally, the video provided unrealistic advice. (R)
After watching the video, people would know exactly what to do.

(R) = item is reverse-scored.

A Likert scale (1: Strongly Disagree to 7: Strongly Agree) is used.

F ADDITIONAL ANALYSES:

Since our sample sizes are smaller, we provide results for non-significant effects so the readers can consider potential false negatives. Note when appropriate, effect sizes are provided throughout to guide the interpretation as they are independent of sample size [84].

F.1 Study 1

Re: Predictors of GPP (Personality traits & Content Appraisal variables): the below table shows the excluded variables in the regression analysis.

Excluded Variables			
	Beta	In t	Sig.
Extraversion	0.095	1.38912	0.169
Agreeable	0.118	1.76777	0.081
Conscientious	-0.034	-0.4866	0.628
Openness	0.012	0.18103	0.857
Information Clarity	-0.024	-0.2655	0.791
Information Importance	0.116	1.31965	0.191

Figure 11: Excluded variables in regression analysis exploring personality traits variables as well as content appraisal variables as predictors of GPP.

Personality Traits & Negative Affect: the below table is the correlation Matrix showing the correlations between personality traits and negative affect.

		Extraversion	Agreeable	Conscientious	Neurotic	Openness	Negative Affect
Extraversion	Correlation Coefficient	1	0.13069	0.042688207	-.234*	0.004796044	-0.163464157
	Sig. (2-tailed)		0.254085	0.710567239	0.039031982	0.966758678	0.152715171
	N	78	78	78	78	78	78
Agreeable	Correlation Coefficient	0.13068967	1	0.169148333	-.276*	0.021052953	-0.170454635
	Sig. (2-tailed)	0.25408461		0.138753383	0.014272173	0.854834928	0.135685443
	N	78	78	78	78	78	78
Conscientious	Correlation Coefficient	0.04268821	0.169148	1	-.398**	0.049577655	-.363**
	Sig. (2-tailed)	0.71056724	0.138753		0.000304695	0.666429123	0.001097303
	N	78	78	78	78	78	78
Neurotic	Correlation Coefficient	-.234*	-.276*	-.398**	1	-0.139575989	.594**
	Sig. (2-tailed)	0.03903198	0.014272	0.000304695		0.222929266	9.9648E-09
	N	78	78	78	78	78	78
Openness	Correlation Coefficient	0.00479604	0.021053	0.049577655	-0.139575989	1	-0.168304717
	Sig. (2-tailed)	0.96675868	0.854835	0.666429123	0.222929266		0.140762323
	N	78	78	78	78	78	78
Negative Affect	Correlation Coefficient	-0.16346416	-0.170455	-.363**	.594**	-0.168304717	1
	Sig. (2-tailed)	0.15271517	0.135685	0.001097303	9.96E-09	0.140762323	
	N	78	78	78	78	78	78

*. Correlation is significant at the 0.05 level (2-tailed).

** . Correlation is significant at the 0.01 level (2-tailed).

Figure 12: Correlations between personality traits and negative affect

F.2 Study 2

Re: Perceived actionability (section 4.5.4) : below, we list results of all pairwise conditions comparison using Mann-Whitney U test (including null effects).

- No solutions condition ($n = 25$; Mean ranking was 22.70) vs. General solutions condition ($n = 29$; Mean ranking was 31.64).
 $U = 242.5$, $Z = -2.09$, $p = .04$ (With Holm-Bonferroni, the alpha was set at .025.), $r = .28$, a small effect.
- General solutions condition ($n = 29$; Mean ranking was 25.71) vs. Actionable solutions condition ($n = 28$; Mean ranking was 32.41).
 $U = 310.50$, $Z = -1.53$, $p = .13$ (With Holm-Bonferroni, the alpha was set at .05), $r = .20$, a small effect.
- No solutions condition ($n = 25$; Mean ranking was 19.88) vs. Actionable solutions condition ($n = 28$; Mean ranking was 33.36).
 $U = 172.00$, $Z = -3.18$, $p = .001^*$ (With Holm-Bonferroni, the alpha was set at .0167), $r = .44$, a medium effect.

Re: Usefulness (section 4.5.4) : below, we list the results of all pairwise comparisons between our video conditions in terms of perceived usefulness using Mann-Whitney U tests.

- No solutions condition ($n = 25$; Mean ranking was 23.98) vs. General solutions condition ($n = 29$; Mean ranking was 30.53).
 $U = 274.50$, $Z = -1.578$, $p = .115$ (With Holm-Bonferroni, the alpha was set at .025), $r = .21$ which is a a small effect .
- General solutions condition ($n = 29$; Mean ranking was 26.19) vs. Actionable solutions condition ($n = 28$; Mean ranking was 31.91).
 $U = 324.50$, $Z = -1.387$, $p = .166$ (With Holm-Bonferroni, the alpha was set at .05), $r = .18$ which is a a small effect .
- No solutions condition ($n = 25$; Mean ranking was 21.52) vs. Actionable solutions condition ($n = 28$; Mean ranking was 31.89).
 $U = 213.00$, $Z = -2.555$, $p = .011^*$ (With Holm-Bonferroni, the alpha was set at .0167), $r = .35$, a medium effect.

Re: Actionability, Anxiety, & GPP (section 4.5.4) : below, we list the results of all pairwise comparisons in reducing the anxiety level using Mann-Whitney U test.

- No solutions condition ($n = 25$; Mean ranking was 23.50) vs. General solutions condition ($n = 29$; Mean ranking was 30.95).
 $U = 262.50$, $Z = -2.252$, $p = .024$ (With Holm-Bonferroni, the alpha was set at .025), $r = .31$ which is a medium effect .
- General solutions condition ($n = 29$; Mean ranking was 28.71) vs. Actionable solutions condition ($n = 28$; Mean ranking was 29.30).
 $U = 397.50$, $Z = -.14$, $p = .89$ (With Holm-Bonferroni, the alpha was set at .05), $r = .02$ which is a small effect .
- No solutions condition ($n = 25$; Mean ranking was 22.00) vs. Actionable solutions condition ($n = 28$; Mean ranking was 31.46).
 $U = 225.00$, $Z = -2.872$, $p = .004$ (With Holm-Bonferroni, the alpha was set at .0167), $r = .39$ which is a medium effect .

Re: Personality traits and Negative Affect (section 4.5.4): Although the exploration of personality traits was not central to Study 2 as it focused on the investigation of the solution for information usefulness, below we provide additional analyses we conducted.

We explored whether personality traits could predict experienced negative affects with linear regression with personality traits as predictors, using the stepwise method. Only neuroticism ($\beta = .37$; $p = .001$) and conscientiousness ($\beta = -.34$; $p < .001$) and remained in the model ($R^2_{adj} = .36$, $p < .001$). Please see below for the excluded variables.

Excluded Variables			
	Beta In	t	Sig.
Extroversion	.108c	1.304	0.195
Agreeable	.042c	0.455	0.650
Openness	.004c	0.045	0.964

Figure 13: Excluded variables in regression analysis exploring personality traits as predictors of experienced levels of negative affect.